arXiv:1208.0294v1 [gr-qc] 1 Aug 2012

CONSTRUCTIVE PROOF OF THE KERR-NEWMAN BLACK HOLE UNIQUENESS: DERIVATION OF THE FULL SOLUTION FROM SCRATCH

REINHARD MEINEL and RENÉ RICHTER

University of Jena, Theoretisch-Physikalisches Institut,
Max-Wien-Platz 1, 07743 Jena, Germany
meinel@tpi.uni-jena.de

The Kerr-Newman black hole solution can be constructed straightforwardly as the unique solution to the boundary value problem of the Einstein-Maxwell equations corresponding to an asymptotically flat, stationary and axisymmetric electro-vacuum spacetime surrounding a connected Killing horizon.

1. Introduction

It was shown in Ref. 1 that the complex Ernst potentials[2] on the symmetry axis of an asymptotically flat, stationary and axisymmetric electro-vacuum spacetime surrounding a connected Killing horizon (including the case of a degenerate horizon) can be constructed uniquely by means of the inverse scattering method.[3,4] This provides a new proof of the Kerr-Newman black hole uniqueness since the full solution can straightforwardly be derived from the axis data, cf. Ref. 5. Here we present a particular method for obtaining the Ernst potentials off the axis, see Sec. 4.

2. Ernst equations and Linear Problem

The stationary and axisymmetric Einstein-Maxwell vacuum equations are equivalent to the Ernst equations[2]

$$f\,\Delta\,\mathcal{E} = (\nabla\mathcal{E} + 2\bar{\Phi}\nabla\Phi)\cdot\nabla\mathcal{E}\,, \quad f\,\Delta\,\Phi = (\nabla\mathcal{E} + 2\bar{\Phi}\nabla\Phi)\cdot\nabla\Phi \tag{1}$$

$$\text{with} \quad f \equiv \Re\,\mathcal{E} + |\Phi|^2\,, \quad \Delta = \frac{\partial^2}{\partial\rho^2} + \frac{1}{\rho}\frac{\partial}{\partial\rho} + \frac{\partial^2}{\partial\zeta^2}\,, \quad \nabla = (\frac{\partial}{\partial\rho}, \frac{\partial}{\partial\zeta})\,, \tag{2}$$

where ρ and ζ are Weyl coordinates and a bar denotes complex conjugation. These equations are the integrability condition of a related Linear Problem (LP):[6,7]

$$\mathbf{Y}_{,z} = \left[\begin{pmatrix} B_1 & 0 & C_1 \\ 0 & A_1 & 0 \\ D_1 & 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & B_1 & 0 \\ A_1 & 0 & -C_1 \\ 0 & D_1 & 0 \end{pmatrix}\right]\mathbf{Y}\,, \tag{3}$$

$$\mathbf{Y}_{,\bar{z}} = \left[\begin{pmatrix} B_2 & 0 & C_2 \\ 0 & A_2 & 0 \\ D_2 & 0 & 0 \end{pmatrix} + \frac{1}{\lambda} \begin{pmatrix} 0 & B_2 & 0 \\ A_2 & 0 & -C_2 \\ 0 & D_2 & 0 \end{pmatrix}\right]\mathbf{Y} \tag{4}$$

$$\text{with} \quad \lambda = \sqrt{\frac{K - \mathrm{i}\bar{z}}{K + \mathrm{i}z}}\,, \quad z = \rho + \mathrm{i}\zeta\,, \quad \bar{z} = \rho - \mathrm{i}\zeta\,, \tag{5}$$

$$A_1 = \bar{B}_2 = \frac{\mathcal{E}_{,z} + 2\bar{\Phi}\Phi_{,z}}{2f}\,, \quad A_2 = \bar{B}_1 = \frac{\mathcal{E}_{,\bar{z}} + 2\bar{\Phi}\Phi_{,\bar{z}}}{2f}\,, \tag{6}$$

$$C_1 = f\bar{D}_2 = \Phi_{,z}\,, \quad C_2 = f\bar{D}_1 = \Phi_{,\bar{z}}\,, \tag{7}$$

where K is the complex “spectral parameter”.

3. Solution on the axis

On the upper part of the symmetry axis ($\rho = 0$, $\zeta > l$), denoted by $\mathcal{A}^+$, a suitably normalized solution to the LP for the black hole problem (horizon at $\rho = 0$, $|\zeta| \le l$) reads[1]

$$\mathcal{A}^+: \quad \mathbf{Y}_+ = \begin{pmatrix} \bar{\mathcal{E}}_+ + 2|\Phi_+|^2 & 1 & \Phi_+ \\ \mathcal{E}_+ & -1 & -\Phi_+ \\ 2\bar{\Phi}_+ & 0 & 1 \end{pmatrix} \begin{pmatrix} F & 0 & 0 \\ G & 1 & L \\ H & 0 & 1 \end{pmatrix}, \tag{8}$$

where we have assumed $\lambda = +1$ for $K \ne \zeta$. The Ernst potentials $\mathcal{E}_+(\zeta)$ and $\Phi_+(\zeta)$ on $\mathcal{A}^+$ are given by

$$\mathcal{E}_+ = 1 - \frac{2M}{\zeta + M - \mathrm{i}J/M}\,, \quad \Phi_+ = \frac{Q}{\zeta + M - \mathrm{i}J/M} \tag{9}$$

together with the parameter relation

$$\frac{l^2}{M^2} + \frac{Q^2}{M^2} + \frac{J^2}{M^4} = 1\,, \tag{10}$$

where M, J and Q denote mass, angular momentum and charge. The functions $F(K)$, $G(K)$, $H(K)$ and $L(K)$ are given by

$$F = \frac{(K - L_1)(K - L_2)}{(K - K_1)(K - K_2)}\,, \quad G = \frac{Q^2 - 2\mathrm{i}J}{(K - K_1)(K - K_2)}\,, \tag{11}$$

$$H = -\frac{2Q(K - L_1)}{(K - K_1)(K - K_2)}\,, \quad L = -\frac{Q}{K - L_1} \tag{12}$$

$$\text{with} \quad L_{1/2} = -M \pm \mathrm{i}\frac{J}{M}\,, \quad K_{1/2} = \pm\sqrt{M^2 - Q^2 - \frac{J^2}{M^2}}\,. \tag{13}$$

4. Solution off the axis

Using the relation[1]

$$\mathbf{Y}(\rho, \zeta, -\lambda) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mathbf{Y}(\rho, \zeta, \lambda) \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{14}$$

together with

$$\mathbf{Y}(\rho, \zeta, 1) = \begin{pmatrix} \bar{\mathcal{E}} + 2|\Phi|^2 & 1 & \Phi \\ \mathcal{E} & -1 & -\Phi \\ 2\bar{\Phi} & 0 & 1 \end{pmatrix} \tag{15}$$

we are led from (8) to the following ansatz for $\mathbf{Y}(\rho,\zeta,\lambda)$:

$$\mathbf{Y}(\rho,\zeta,\lambda)=\begin{pmatrix}\psi(\rho,\zeta,\lambda) & \psi(\rho,\zeta,-\lambda) & \alpha(\rho,\zeta,\lambda)\\ \chi(\rho,\zeta,\lambda) & -\chi(\rho,\zeta,-\lambda) & \beta(\rho,\zeta,\lambda)\\ \varphi(\rho,\zeta,\lambda) & \varphi(\rho,\zeta,-\lambda) & \gamma(\rho,\zeta,\lambda)\end{pmatrix} \tag{16}$$

with

$$\psi=1+a_1\left(\frac{1}{\kappa_1-\lambda}-\frac{1}{\kappa_1+1}\right)+a_2\left(\frac{1}{\kappa_2-\lambda}-\frac{1}{\kappa_2+1}\right), \tag{17}$$

$$\chi=1+b_1\left(\frac{1}{\kappa_1-\lambda}-\frac{1}{\kappa_1+1}\right)+b_2\left(\frac{1}{\kappa_2-\lambda}-\frac{1}{\kappa_2+1}\right), \tag{18}$$

$$\varphi=c_1\left(\frac{1}{\kappa_1-\lambda}-\frac{1}{\kappa_1+1}\right)+c_2\left(\frac{1}{\kappa_2-\lambda}-\frac{1}{\kappa_2+1}\right), \tag{19}$$

$$\alpha=\Phi+\frac{\alpha_0}{K-L_1}\,,\quad \beta=-\Phi\,\frac{\lambda(K+\mathrm{i}z)}{K-L_1}\,,\quad \gamma=1+\frac{\gamma_0}{K-L_1}\,, \tag{20}$$

$$\text{where}\quad \kappa_i=\sqrt{\frac{K_i-\mathrm{i}\bar{z}}{K_i+\mathrm{i}z}}\quad (\mathcal{A}^+:\quad \kappa_i=+1)\,. \tag{21}$$

A careful discussion of the regularity conditions at

$$\lambda_i=\sqrt{\frac{L_i-\mathrm{i}\bar{z}}{L_i+\mathrm{i}z}} \tag{22}$$

leads to a set of linear algebraic equations which uniquely determine the unknowns $a_i(\rho,\zeta)$, $b_i(\rho,\zeta)$, $c_i(\rho,\zeta)$, $\alpha_0(\rho,\zeta)$, $\gamma_0(\rho,\zeta)$ and $\Phi(\rho,\zeta)$. According to (15), $\mathcal{E}(\rho,\zeta)$ is given by $\chi(\rho,\zeta,1)$. The resulting Ernst potentials are the well-known expressions

$$\mathcal{E}=1-\frac{2M}{r-\mathrm{i}(J/M)\cos\theta}\,,\quad \Phi=\frac{Q}{r-\mathrm{i}(J/M)\cos\theta} \tag{23}$$

with Boyer-Lindquist coordinates r and θ related to our Weyl coordinates ρ and ζ by

$$\rho=\sqrt{r^2-2Mr+J^2/M^2+Q^2}\,\sin\theta\,,\quad \zeta=(r-M)\cos\theta\,. \tag{24}$$

References

1. R. Meinel, *Class. Quantum Grav.* **29**, 035004 (2012), arXiv:1108.4854.
2. F. J. Ernst, *Phys. Rev.* **168**, 1415 (1968).
3. V. Belinski and E. Verdaguer, *Gravitational Solitons* (Cambridge University Press, Cambridge, 2001).
4. R. Meinel, M. Ansorg, A. Kleinwächter, G. Neugebauer and D. Petroff, *Relativistic Figures of Equilibrium* (Cambridge University Press, Cambridge, 2008).
5. I. Hauser and F. J. Ernst, *J. Math. Phys.* **22**, 1051 (1981).
6. G. A. Alekseev, *JETP Lett.* **32**, 277 (1980).
7. G. Neugebauer and D. Kramer, *J. Phys. A: Math. Gen.* **16**, 1927 (1983).